

OFFICE OF NAVAL RESEARCH

CONTRACT N00014-97-1-0066

R&T Code 33e 1806

Dr. Judah Goldwasser

Technical Report No. 102

COMPUTED HEATS OF FORMATION

by

Peter Politzer, M. Edward Grice, Monica C. Concha and Pat Lane

Department of Chemistry  
University of New Orleans  
New Orleans, LA 70148

April 9, 1998

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.









DTIC QUALITY INSPECTED 2

19980429 000

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.







1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE April 9, 1998	3. REPORT TYPE AND DATES COVERED Technical
4. TITLE AND SUBTITLE Computed Heats of Formation			5. FUNDING NUMBERS N00014-97-1-0066 Dr. Judah Goldwasser R&T Code 33e 1806
6. AUTHOR(S) Peter Politzer, M. Edward Grice, Monica C. Concha and Pat Lane			8. PERFORMING ORGANIZATION REPORT NUMBER 102
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of New Orleans Department of Chemistry New Orleans, Louisiana 70148			10. SPONSORING / MONITORING AGENCY REPORT NUMBER
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Code 333 800 N. Quincy Street Arlington, VA 22217			11. SUPPLEMENTARY NOTES
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release. Unlimited distribution.			12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 words) Computed heats of formation for 1 - 8. <div style="display: flex; justify-content: space-around; align-items: flex-start;"><div style="text-align: center;"> 1</div><div style="text-align: center;"> 2</div><div style="text-align: center;"> 3: Z = H 4: Z = NO<sub>2</sub> 5: Z = NF<sub>2</sub></div><div style="text-align: center;"> 6</div></div> <div style="display: flex; justify-content: space-around; align-items: flex-start; margin-top: 20px;"><div style="text-align: center;"> 7</div><div style="text-align: center;"> 8</div><div style="text-align: center;"> 1: <math>\Delta H_f^{298K}</math> (solid) = 231 cal/g 2: <math>\Delta H_f^{298K}</math> (solid) = 491 cal/g 3: <math>\Delta H_f^{298K}</math> (solid) = 150 cal/g 4: <math>\Delta H_f^{298K}</math> (solid) = 157 cal/g</div><div style="text-align: center;"> 5: <math>\Delta H_f^{298K}</math> (solid) = 132 cal/g 6: <math>\Delta H_f^{298K}</math> (solid) = 235 cal/g 7: <math>\Delta H_f^{298K}</math> (solid) = 1.5 cal/g 8: <math>\Delta H_f^{298K}</math> (solid) = -584 cal/g</div></div>			

We have computed heats of formation for compounds **1 - 8** (Table 1). The first five are target compounds proposed by M. Trudell (University of New Orleans); **6 - 8** have recently been prepared by R. Schmitt and J. Bottaro (SRI). For the molecular systems **1 - 7**, we used our density functional procedure to obtain gas phase heats of formation, which were converted to liquid and solid state values by subtracting, respectively, the heats of vaporization and sublimation. The latter are determined by means of relationships that we have developed involving the computed electrostatic potential on the molecular surface [2,3]. (Vibrational energies were obtained from the molecular stoichiometries [4].) For the ionic solid **8**, the heat of formation was calculated using the lattice enthalpy and the gas phase heats of formation of the positive and negative ions; the lattice enthalpy was computed from our recently-developed relationship involving anionic surface electrostatic potentials [5]. For comparison, the experimental solid phase heats of formation of HMX and RDX are, respectively, 60.4 cal/g and 76.1 cal/g [6].

#### References:


1. D. Habibollahzadeh, M. E. Grice, M. C. Concha, J. S. Murray and P. Politzer, *J. Comp. Chem.* **16**, 654 (1995).
2. J. S. Murray and P. Politzer, *Quantitative Treatments of Solute/Solvent Interactions*, J.S. Murray and P. Politzer, Eds., Elsevier, Amsterdam, 1994, ch. 8.
3. P. Politzer, J. S. Murray, M. E. Grice, M. DeSalvo and E. Miller, *Mol. Phys.* **91**, 923 (1997).
4. M. E. Grice and P. Politzer, *Chem. Phys. Lett.* **244**, 295 (1995).
5. P. Politzer and J. S. Murray, *J. Phys. Chem. A*, **102**, 1018 (1998).
6. J. Köhler and R. Meyer, *Explosives*, 4th ed., VCH Publishers, New York, 1993.

Table 1. Computed heats of formation.

1		$\Delta H_f^{298K}(\text{gas}) = 73.6 \text{ kcal/mole} = 341 \text{ cal/g}$ $\Delta H_f^{298K}(\text{liquid}) = 60.0 \text{ kcal/mole} = 278 \text{ cal/g}$ $\Delta H_f^{298K}(\text{solid}) = 49.8 \text{ kcal/mole} = 231 \text{ cal/g}$
2		$\Delta H_f^{298K}(\text{gas}) = 155 \text{ kcal/mole} = 604 \text{ cal/g}$ $\Delta H_f^{298K}(\text{liquid}) = 139 \text{ kcal/mole} = 544 \text{ cal/g}$ $\Delta H_f^{298K}(\text{solid}) = 126 \text{ kcal/mole} = 491 \text{ cal/g}$
3		$\Delta H_f^{298K}(\text{gas}) = 80.6 \text{ kcal/mole} = 263 \text{ cal/g}$ $\Delta H_f^{298K}(\text{liquid}) = 64.3 \text{ kcal/mole} = 210 \text{ cal/g}$ $\Delta H_f^{298K}(\text{solid}) = 45.8 \text{ kcal/mole} = 150 \text{ cal/g}$
4		$\Delta H_f^{298K}(\text{gas}) = 104 \text{ kcal/mole} = 264 \text{ cal/g}$ $\Delta H_f^{298K}(\text{liquid}) = 87.5 \text{ kcal/mole} = 221 \text{ cal/g}$ $\Delta H_f^{298K}(\text{solid}) = 62.2 \text{ kcal/mole} = 157 \text{ cal/g}$
5		$\Delta H_f^{298K}(\text{gas}) = 96.5 \text{ kcal/mole} = 236 \text{ cal/g}$ $\Delta H_f^{298K}(\text{liquid}) = 79.1 \text{ kcal/mole} = 194 \text{ cal/g}$ $\Delta H_f^{298K}(\text{solid}) = 54.0 \text{ kcal/mole} = 132 \text{ cal/g}$
6		$\Delta H_f^{298K}(\text{gas}) = 108 \text{ kcal/mole} = 370 \text{ cal/g}$ $\Delta H_f^{298K}(\text{liquid}) = 91.1 \text{ kcal/mole} = 312 \text{ cal/g}$ $\Delta H_f^{298K}(\text{solid}) = 68.7 \text{ kcal/mole} = 235 \text{ cal/g}$

(continued)

Table 1. Computed heats of formation (continued).

7		$\Delta H_f^{298K}(\text{gas}) = 20.5 \text{ kcal/mole} = 124 \text{ cal/g}$ $\Delta H_f^{298K}(\text{liquid}) = 8.23 \text{ kcal/mole} = 49.8 \text{ cal/g}$ $\Delta H_f^{298K}(\text{solid}) = 0.24 \text{ kcal/mole} = 1.5 \text{ cal/g}$
8	$\text{K}^+ \left[ \text{N} \begin{array}{l} \nearrow \text{F} \\ \searrow \text{NO}_2 \end{array} \right]^-$	$\Delta H_f^{298K}(\text{solid}) = -69.0 \text{ kcal/mole} = -584 \text{ cal/g}$